

Beta-Globin Gene Mutations in India and Their Linkage to β -Haplotypes

Anju Gupta*, Swati Sarwai*, Neelam Pathak** and Sarita Agarwal*

*Department of Genetics, Sanjay Gandhi Post Graduate Institute of Medical Science,
Lucknow 226 014, Uttar Pradesh, India

**Department of Biotechnology, Integral University, Lucknow 226 026, Uttar Pradesh, India

KEYWORDS Haplotypes; β -thalassemia; β -globin gene; Asian Indian; DNA polymorphism; RFLP

ABSTRACT A total of 124 chromosomes of 64 unrelated Indian β -thalassemia and β -thalassemia patients along with their family members were studied for their haplotype pattern and mutations. These included, 35 with β -thalassemia major, 4 with thalassemia trait, and 25 with β -thalassemia. Fourteen mutations were detected by PCR and Sequencing. The most common mutation IVS1-5 (G-C) was linked with 8 different haplotypes. Nineteen haplotypes were found on β -thalassemia mutations, with haplotype (+ - - - + -) being the most widespread and was found associated with 39 chromosomes of IVS1-5 (G-C), 2 of HPFH and 1 each of CD41/42(-CTTT) and CD16 (-C).

INTRODUCTION

The technique of RFLP depends on the property of restriction endonuclease to fragment DNA at specific recognition sites and then electrophoresis to separate the resulting fragment according to their size. Variations from the pattern seen in "normal" DNA samples constitute restriction fragment length polymorphisms or RFLP's.

In β -thalassemia, a heterogeneous genetic disorder of β -globin synthesis, a strong association between particular β -globin gene haplotype and specific mutation was suggested (Orkin et al. 1982). It was only later elucidated that the same haplotype may be associated with different mutations, while the same mutation may be associated with many different haplotypes (Cheng et al. 1984; Orkin and Kazazian 1984; Wong et al. 1986). The aim of this study is to investigate the degree of correlation between different β -globin gene mutations and haplotypes

MATERIALS AND METHODS

Subjects: Sixty-four unrelated Indian families with at least one index case with thalassemia were included in this study. These families were referred to our center from hospitals in 3 different states of India [viz 47 from Uttar Pradesh (U.P.), 7 from West Bengal and 10 from Gujarat].

Methods: Hematological analysis was performed on Sysmex automated cell counter.

HbA₂ was estimated by column chromatography (Schleider et al. 1977) while HbF was determined by alkali denaturation test (Betke et al. 1959) DNA was extracted by the phenol-chloroform method (Poncz et al. 1982). The samples were screened for the 5 common, 5 less common, and few rare mutations reported among Asian Indians (Gupta et al. 2003) first by Reverse dot blot (Gorashaker et al. 1997) and then by Amplification refractory mutation system-PCR (Newton et al. 1989) These samples were then subjected to nested PCR followed by single-stranded conformational polymorphism (SSCP) (Gupta and Agarwal 2003) and sequencing. Haplotype analysis was performed by RFLP-PCR (Sutton et al. 1989). Seven restriction endonuclease sites within the β -globin gene cluster (Fig. 1) were used to construct a haplotype (*Hind*II5'e, *Xmn*I5'Gg, *Hind*III Gg, *Hind*III Ag, *Hinc*II5'jb, *Hinf*I5'b, and *Ava*IIb).

DNA was amplified using 15.0pM of each primer, 2.5mM of MgCl₂ (Bangalore Genei), 10.0mM of dNTPs (Bangalore Genei) and 1 Unit of *taq* polymerase (Bangalore Genei). This sample mix was subjected to hot start consisting of denaturation at 94°C for 7 min, annealing at 65°C for 1 min and extension at 72°C for 1.30 min and finally to 24 PCR cycles. Each cycle was 94°C for 1 min, 1 min annealing at 65°C and an extension of 1.30 min at 72°C (Fig. 2).

The digestion reaction was performed in an overnight incubation of 10mg of each PCR product with the correspondent endonuclease according to the manufacturer recommendation.




Fig. 1. Diagrammatic representation of 7 RFLP sites in β -globin gene cluster.




Fig. 2. Amplicons for different RFLP sites in β -globin gene cluster. Left to right marker, Xmn I, Hind III G, 3'Hinc II $\gamma\beta$, Hinf I, 5'Hind III $\phi\beta$, Ava II, 5' Hinc IIE

1	2	3	4	5	6	7	8	9	10	11	12	13
	HindII			HindII		M		AvaII			HindIII	
	U 760			794				328			635	
	D 446, 314			690, 104				228, 110			327, 308	



Fig. 3. Gel photograph showing PCR products after digestion with respective enzymes. 1,2,3 - HindII site absent (-/-), 4,5-(+/-), 6-(+/+), 8, 9, 10- (+/+), 11, 12, 13-(-/-).

The digestion profiles were obtained by agarose gel electrophoresis with subsequent staining with ethidium bromide and visualization in a UV transilluminator (Fig. 3).

RESULTS

Characterization of Mutations

Of the 124 chromosomes studied, 72 were of IVS1-5(G-C), 6 of IVS1-1(G-T), 5 of CD41/42(-CTTT), 3 of CD8/9(+G) and 2 each of CD15(-T), CD16(-C) and HPFH, 1 each of 619 bp deletion, CD15(G-A), -29(A-G), CD 30(G-C), Ini (ATG-ACG), a novel 8bp deletion, a 17 bp deletion at CD126-131 and 25 of CD26.

The five most common Asian Indian mutations accounted for 70% of the total allele with IVS1-5 (G-C) being the predominant mutation (58%).

In 10 β -thalassemia chromosomes and 6 HbE chromosomes, haplotypes could not be assigned because of heterozygosity at one or more polymorphic site and nonavailability of family members.

Haplotype Study

A total of 19 haplotypes were found on β -thalassemia chromosomes (Table 1). The most common mutation IVS1-5 (G-C) was associated with 8 different haplotypes with haplotype {+ - - - - +} being the predominant (54%). It was found on 32 chromosomes of UP origin, 6 of W. Bengal and 1 of Gujarat. This haplotype was also associated with CD16 (-C) and CD41/42 (-CTTT). The second haplotype associated with the mutation was {+ - - - - +} that was found in 8 chromosomes (1 U.P. and 7 Gujarati). The third most common haplotype associated with IVS1-5 was {- - + + - -} and was present on 5 chromosomes. The 3 major haplotypes associated with IVS1-5 accounted for a total of 72.2%.

IVS1-1 (G-T) mutation was found on 2 haplotypes, while CD8/9 (+G) and CD41/42 (-CTTT) were found on 3 haplotypes each. IVS1-1 (G-T) and CD8/9 (+G) are present on the same chromosomal background. CD15 (G-A), CD16 (-C), 619bp del, HPFH and CD26 were found on the same haplotype as IVS1-5 (G-C). CD30 (G-C) and novel 8bp deletion shared the same haplotype. Ini (ATG-ACG), 17 bp del, CD15 (-T)

Table 1: Association of β -thalassemia/HbE mutations and haplotypes.

Mutations	Haplotype	U.P.	W.Bengal	Gujarat
IVS1-5(G-C)	+ - - - - +	32	6	1
	- - + + - -	5	-	
	- - - - - +	3	-	1
	+ - - - - -			2
	+ - - - - +	1		7
	+ - - - - +	1		
	- - + - - +			1
	- + - - - +			1
IVS1-1(G-T)	- + + - - +	4		
	- + + + + +	2		
CD8/9(+G)	+ + + - - -			1
	- + + - - +	1		
	- + + + + +	1		
CD15 (-T)	- - + + - -	2		
CD15 (G-A)	+ - - - - +			1
-29 (A-G)	+ + + - - +	1		
CD30 (G-C)	+ + + + + +	1		
CD16 (-C)	+ - - - - -	1		
	+ - - - - +	1		
CD41/42 (-CTTT)	- - - - - +			2
	+ - - - - -			1
	- - + - - +	2		
Ini (ATG-ACG)	- - + + + +	1		
-8bp del	+ + + + + +	1		
-17bp del	- - + + - -	1		
-619bp del	+ - - - - -			1
HPFH	+ - - - - -	2		
CD26	- + - - - +	1	3	
	+ - - - - +	13	0	
Normal	+ - - - - +	41		
	+ + - - - -	12		
	- + + - - +	1		
	- + + - - -	2		
	- - + - - +	4		

and -29 (A-G) were found on different chromosomal backgrounds. Both the chromosomes with CD15 (-T), a rare mutation in Indians was found to have the same haplotype pattern.

A total of 25 b^E chromosomes were analyzed for the β -globin gene cluster haplotypes (18 of U.P. and 7 of W. Bengal origin). Strong linkage of b^E mutation with one particular haplotype {+ - - - - +} in 13 chromosomes (72.2%) of U.P. origin was observed. However, in the W. Bengal group haplotype {- + - - - +} significantly contributed to linkage with the b^E mutation (42.8%)

Among the 73 normal chromosomes, 5 different β -globin cluster haplotypes were found. With the exception of one {+ - - - - +}, none of the β -thalassemia chromosomes were represented among the normal haplotypes. The predominant haplotype {+ - - - - +} was found associated with IVS1-5 (G-C) but with lower frequency.

DISCUSSION

Analysis of DNA polymorphic sites is a valuable tool in relating human genetic variation to disease and for understanding human evolutionary history. The beta-globin locus contains several single-base restriction fragment length polymorphism (RFLP) sites throughout chromosome 11. In addition to these polymorphic sequence repeats, others are being studied in order to expand our knowledge concerning the role between haplotype-genotype and phenotype associations (eg., presence of XmnI leads to elevation of HbF and thereafter buffering the severity of the disease).

β -thalassemia alleles found among our patients are typical of Asian Indians. Our results reveal that the five most common mutations are IVS1-5(G-C), IVS1-1(G-T), CD8/9(+G), CD41/42(-CTTT) and -619bp constitute 70.0% of the allele.

The IVS1-5 is the most frequently observed mutation present in β -thalassemic chromosomes represents 58.0% of the 124 β -thalassemic chromosomes studied.

It has been reported in high frequency in Indonesians and Melanesians but in low frequency in Chinese and Thais. In Indonesians and Melanesians this variant is mostly associated with the haplotype {+-----++}(Lie-Injo et al. 1989), in Chinese with haplotype {------++++}. In our case we found that eight different haplotype harbored IVS1-5 (G-C) mutation, with {+-----} being the most common. There are two plausible explanations for such extreme heterogeneity associated with this mutation. (1) The mutation arose more than once in the population, the chances of which are very rare. (2) multiple haplotypes are result of simple recombination events, which is more likely as 5' region to beta-globin gene is a hotspot for recombination. The different and vast number of haplotypes associated with this mutation, its high frequency and omnipresence is suggestive its independent as well as most ancient origin among Asian Indians.

CD 41/42 is another prevalent mutation found in Southeast Asian countries. This mutation was first reported in Taiwanese (Kimura et al. 1983), later in Chinese (Wong et al. 1986) and in Asian Indians (Kazazian et al. 1984). Its widespread distribution, high frequency and association with different haplotype and framework suggest a separate/early origin. In our patients it was found on three chromosomes of IVS1-5. Unlike what

was reported by Varawalla et al. (1992) and Thein et al. (1984), in our patient group -619 bp, CD8/9 and IVS1-1 are not confined to a particular region or group and are not in strong linkage disequilibria with their haplotypes. Infact CD8/9 and IVS1-1 were found on the same chromosomal background. The reason for this could be small sample size. Both the chromosomes with CD15 (-T), having the same haplotype pattern can be explained on the grounds that the couple in which it was found was first cousins. -29 (A-G) mutation is not reported in Indians but has been found in Malay (Tan et al. 2004), Chinese (Mo et al. 2004), American Blacks (Antonarakis et al. 1984), Guadeloupeans (Romana et al. 1996), Moroccans (Lemsaddek et al. 2004), Tunisians (Chouk et al. 2004) and Cubans (Muniz et al. 2000). The mutation was found associated with different haplotypes in different population suggesting an independent origin in those populations.

The high mutational heterogeneity observed in this study has provided useful information about the types and geographic distribution of β -thalassemia mutations in India, thus allowing rapid detection of mutations in couples at risk.

In addition, presence of 19 different haplotype on β -thalassemia chromosomes proves that this genetic diversity has originated from both new mutational events and gene flow due to population migration.

ACKNOWLEDGEMENTS

We would like to thank the Indian Council of Medical Research, Department of Science and Technology, New Delhi, India, for their financial assistance. The authors are also thankful to Ms Vandana Arya for her help.

REFERENCES

- Antonarakis SE, Irkin SH, Cheng TC, Scott AF, Sexton JP, Trusko SP, Charache S, Kazazian HH Jr 1984. Beta-Thalassemia in American Blacks: novel mutations in the "TATA" box and an acceptor splice site. *Proc Natl Acad Sci U S A*, **81**: 1154-1158.
- Betke K, Marti HR, Schlicht I 1959. Estimation of small percentage of fetal hemoglobin. *Nature*, **184**: 1877-1879.
- Chehab FF, Der Kaloustian V, Khouri FP, Deeb SS, Kan YW 1987. The molecular basis of beta-thalassemia in Lebanon: Application to prenatal diagnosis. *Blood*, **69**: 1141-1145.
- Cheng TC, Orkin SH, Antonarakis SE, Potter MJ, Sexton JP, Markham AF, Giardina PJ, Li A, Kazazian HH Jr

1984. Beta-Thalassemia in Chinese: use of *in vivo* RNA analysis and oligonucleotide hybridization in systematic characterization of molecular defects. *Proc Natl Acad Sci U S A*, **81**: 2821-2825.
- Chouk I, Daoud BB, Mellouli F, Bejaoui M, Gerard N, Dellagi K, Abbas S 2004. Contribution to the description of the beta-thalassemia spectrum in Tunisia and the origin of mutation diversity. *Hemoglobin*, **28**: 189-195.
- Gorashakar AC, Pawar AR, Nadkarni AH, Lu CY, Mohanty D, Krishnamoorthy R, Besmond C, Colah RB 1997. Application of covalent reverse dot-blot hybridization for rapid prenatal diagnosis of the common Indian thalassemia syndromes. *Ind J Hematol Blood Transfusion*, **15**: 10-14.
- Gupta A, Agarwal S 2003. Efficiency and cost effectiveness: PAGE-SSCP versus MDE and Phast gels for the identification of unknown beta thalassaemia mutations. *Mol Pathol*, **56**: 237-239.
- Gupta A, Hattori Y, Gupta UR, Sarwai S, Nigam N, Singhal P, Agarwal S 2003. Molecular genetic testing of beta-thalassemia patients of Indian origin and a novel 8-bp deletion mutation at codons 36/37/38/39. *Genet Test*, **7**: 163-168.
- Huang S, Wong C, Antonarakis SE, Ro-lien T, Lo WH, Kazazian HH Jr 1986. The same "TATA" box beta-thalassemia mutation in Chinese and US blacks: another example of independent origins of mutation. *Hum Genet*, **74**: 162-164.
- Kazazian HH Jr, Orkin SH, Antonarakis SE, Sexton JP, Boehm CD, Goff SC, Waber PG 1984. Molecular characterization of seven beta-thalassemia mutations in Asian Indians. *EMBO J*, **3**: 593-596.
- Kimura A, Matsunaga E, Takihara Y, Nakamura T, Takagi Y, Lin S, Lee H 1983. Structural analysis of a beta-thalassemia gene found in Taiwan. *J Biol Chem*, **258**: 2748-2749.
- Lemsaddek W, Picanco I, Seuanes F, Nogueira P, Mahmal L, Benchekroun S, Khattab M, Osorio-Almeida L 2004. The beta-thalassemia mutation/haplotype distribution in the moroccan population. *Hemoglobin*, **28**: 25-37.
- Lie-Injo LE, Cai SP, Wahidijat I, Moeslichan S, Lim ML, Evangelista L, Doherty M, Kan YW 1989. Beta-thalassemia mutations in Indonesia and their linkage to beta haplotypes. *Am J Hum Genet*, **45**: 971-975.
- Mo QH, Zhu H, Li LY, Xu XM 2004. Reliable and high-throughput mutation screening for beta-thalassemia by a single-base extension/fluorescence polarization assay. *Genet Test*, **8**: 257-262.
- Muniz A, Martinez G, Lavinha J, Pacheco P 2000. Beta-thalassaemia in Cubans: Novel allele increases the genetic diversity at the HBB locus in the Caribbean. *Am J Hematol*, **64**: 7-14.
- Newton CR, Graham A, Heptinstall LE, Powell SJ, Summers C, Kalsheker N, Smith JC, Markham AF 1989. Analysis of any point mutation in DNA. The amplification refractory mutation system (ARMS). *Nucleic Acids Res*, **1**: 2503-2516.
- Orkin SH, Kazazian HH Jr 1984. The mutation and polymorphism of the human beta-globin gene and its surrounding DNA. *Annu Rev Genet*, **18**: 131-71.
- Orkin SH, Kazazian HH Jr, Antonarakis SE, Goff SC, Boehm CD, Sexton JP, Waber PG, Giardina PJ 1982. Linkage of beta-thalassaemia mutations and beta-globin gene polymorphisms with DNA polymorphisms in human beta-globin gene cluster. *Nature*, **296**: 627-631.
- Poncz M, Solowiejczyk D, Harpel B, Mory Y, Schwartz E, Surrey S 1982. Construction of human gene libraries from small amounts of peripheral blood: analysis of b-like globin genes. *Hemoglobin*, **6**: 27-35.
- Romana M, Keclard L, Guillemin G, Lavocat E, Saint-Martin C, Berchel C, Merault G 1996. Molecular characterization of beta-thalassemia mutations in Guadeloupe. *Am J Hematol*, **53**: 228-233.
- Schleider CT, Mayson SM, Huisman THJ 1977. Further modification of the microchromatographic determination of hemoglobin A2. *Hemoglobin*, **1**: 503-507.
- Sutton M, Bouhassira EE, Nagel RL 1989. Polymerase chain reaction amplification applied to the determination of β -like globin gene cluster haplotypes. *Am J Hematol*, **32**: 66-69.
- Tan JA, George E, Tan KL, Chow T, Tan PC, Hassan J, Chia P, Subramaniam R, Chandran R, Yap SF 2004. Molecular defects in the beta-globin gene identified in different ethnic groups/populations during prenatal diagnosis for beta-thalassemia: A Malaysian experience. *Clin Exp Med*, **4**: 142-147.
- Thein SL, Old JM, Wainscoat JS, Petrou M, Modell B, Weatherall DJ 1984. Population and genetic studies suggest a single origin for the Indian deletion beta thalassaemia. *Br J Haematol*, **57**: 271-278.
- Thein SL, Winichagoon P, Hesketh C, Best S, Fucharoen S, Wasi P, Weatherall DJ 1990. The molecular basis of beta-thalassemia in Thailand: Application to prenatal diagnosis. *Am J Hum Genet*, **47**: 369-375.
- Varawalla NY, Fitches AC, Old JM 1992. Analysis of beta-globin gene haplotypes in Asian Indians: origin and spread of beta-thalassaemia on the Indian subcontinent. *Hum Genet*, **90**: 443-449.
- Wong C, Antonarakis SE, Goff SC, Orkin SH, Boehm CD, Kazazian HH Jr 1986. On the origin and spread of beta-thalassemia: recurrent observation of four mutations in different ethnic groups. *Proc Natl Acad Sci U S A*, **83**: 6529-6532.