ISSN 0972-3757 International Journal of HUMAN GENETICS



Int J Hum Genet, 17(3): 135-139 (2017) DOI:10.31901/24566330.2017/17.03.06

Brief Communication

Parental Consanguineous Marriages are Associated with Early Age of Onset of Schizophrenia in a Pakistani Cohort

Warda Fatima^{1*}, Saqib Mahmood², Shahida Hasnain^{1,3}, Nusrat Habib Rana⁴, Nasar Sayeed Khan^{5@}, Farooq Naeem^{5%} and Muhammad Ayub^{5#}

¹Department of Microbiology and Molecular Genetics, University of the Punjab, Lahore, Pakistan ²Department of Allied Health Sciences, University of Health Sciences, Lahore, Pakistan ³The Women University Multan, Multan, Pakistan ⁴Punjab Institute of Mental Health, Lahore, Pakistan ⁵Department of Psychiatry, Queen's University, Kingston, Canada

KEYWORDS Age of Onset. Family History. Parental Consanguinity. Schizophrenia

ABSTRACT Schizophrenia is a complex multifactorial psychiatric disorder. Both environmental and genetic factors are thought to play a role in the development of this condition. Many studies have shown an association between consanguinity and schizophrenia. A study from Iran has examined the association between age at onset and consanguinity. In this study, the researchers found a strong relationship between age of onset of schizophrenia symptoms and parental cousin marriages. There was no effect of positive family history of disease on age of onset.