

Full text open access online (Since 2001)



Kamla-Raj IJHG 2024

PRINT: ISSN 0972-3757 ONLINE: ISSN 2456-6330

Int J Hum Genet, 24(1): 44-52 (2024)

DOI: 10.31901/24566322.2024/24.01.875

Common Thrombophilic Mutations among Sickle Cell Disease Patients in the Western Province of Saudi Arabia

**Rasha Desuqi¹, Najlaa Filimban¹, Raed Alserihi¹, Anwar Refaei¹,
Yahya Almotawif², Nora Hakami¹ and Talal Qadah¹**

*¹Department of Medical Laboratory Sciences, Faculty of Applied Medical Sciences,
King Abdulaziz University, Jeddah, Saudi Arabia*

*²Department of Medical Laboratory Technology, College of Applied Medical Sciences,
Taibah University, Al-Madinah Al-Munawwarah Saudi Arabia*

KEYWORDS Thrombophilia. Sickle Cell Disease. Thrombosis Risk. Genetic Screening. Vascular Complication

ABSTRACT Sickle cell disease (SCD), linked to vascular thrombosis, is an autosomal recessive disorder. Among the various thrombophilic mutations, Factor V Leiden (FVL) G1691A, prothrombin (PRT) G20210A, and methylenetetrahydrofolate reductase (MTHFR) C677T stand out as genetic alterations linked to thrombotic consequences. Thus, the study aims to examine the prevalence of these mutations in SCD patients, utilizing the polymerase chain reaction-based technique. The study design is an observational study. The study results revealed that there are significant levels of the heterozygous form of FVL G1691A and PRT G20210A in the SCD patient population compared to control subjects. While, MTHFR C677T showed no statistical significance. None of the homozygous forms of the three mutations was statistically significant. However, the incidence of FVL, PRT, and MTHFR mutations was higher among female participants than male participants in the study. The high prevalence of these mutations suggests that they may be significant risk factors for the vascular complications in this population. However, further studies are required to validate the current outcomes.