



SETD1B Inhibits the Progression of Diffuse Large B Cell Lymphomas Via Regulating RAB 17

Jingwen Gu¹, Tengfei Long², Shan Gao³ and Tianping Chen^{4,*}

^{1,3,4}*Department of Hematology and Oncology, Anhui Provincial Children's Hospital, Hefei 230022, China*

²*Hefei Ion Medical Center, the First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei 230031, China*

KEYWORDS Apoptosis. DLBCL. Cancer. Histone Methylation. SET Domain Containing 1B

ABSTRACT The purpose of this study is to explore the role of SET domain containing 1B (SETD1B) in diffuse large B cell lymphomas (DLBCL). Gene expression is analysed by quantitative reverse transcription PCR and Western blot. Cell behaviours are analysed by cell counting kit-8, 5-Ethynyl-2'-deoxyuridine, terminal deoxynucleotidyl transferase dUTP risk end labelling, and flow cytometry assays. The interaction between SETD1B and RAB family member 17 (RAB17) is analysed by co-immunoprecipitation and chromatin immunoprecipitation assays. The researchers found that SETD1B expression is downregulated in DLBCL. Overexpression SETD1B promotes the apoptosis of DLBCL cells. Moreover, SETD1B promotes the H3K4me3 methylation and upregulation of RAB17. However, RAB17 knockdown promotes the survival of DLBCL cells. In summary, SETD1B exerts anti-tumour function in DLBCL. SETD1B drive the apoptosis of DLBCL via mediating histone methylation of RAB17.

INTRODUCTION

Diffuse large B-cell lymphoma (DLBCL) is a popular hematological tumor (Kotlov et al. 2021; Nakhoda et al. 2023; Schmitz et al. 2018). DLBCL is characterised by the rapid proliferation of B cells. Cancerous B cells in DLBCL lose their ability to function properly and can spread to various organs (Chen et al. 2022; Pi et al. 2022). DLBCL is a highly heterogeneous disease, with significant variations in clinical presentation, genetic features, and treatment outcomes (Chapuy et al. 2018). The World Health Organisation (WHO) classification system recognises over a dozen subtypes of DLBCL, each with distinct genetic and molecular characteristics (Dabrowska-Iwanicka and Nowakowski 2024). Understanding the underlying molecular mechanisms driving DLBCL is crucial for developing more effective targeted therapies and improving patient outcomes.

Objectives

The researchers aimed to explore the role of SETD1B in DLBCL and the underlying molecular mechanisms.

Recent Literature Review

SET domain containing 1B (SETD1B) is a histone lysine methyltransferase (Kranz and Anastassiadis 2020). SETD1B modifies histones by adding methyl groups, thereby influencing gene expression and cellular function (Roston et al. 2021). Recent studies have identified SETD1B mutations and deletions in a significant proportion of DLBCL cases. Specifically, SETD1B alterations occur in approximately 16 percent of DLBCL patients. Loss-function mutation of SETD1B is associated with resistance to apoptosis of cancer cells. SETD1B deficiency has been shown to confer resistance to Venetoclax, highlighting its importance in the pathogenesis of DLBCL (Portelinha et al. 2024). Its loss leads to the downregulation of these proteins, promoting cell survival and contributing to lymphomagenesis (Ozyerli-Goknar et al. 2023). Additionally, SETD1B cooperates with other genetic abnormalities, such as the loss of lysine methyltransferase 2D, to drive lymphoma de-

*Address for correspondence:

Tianping Chen,
Department of Hematology and Oncology,
Anhui Provincial Children's Hospital,
39# Wangjiang East Road, Hefei 230022, China
Tel: 0551-63293091
E-mail: tianping_chen@outlook.com

velopment (Shimada et al. 2021). Notwithstanding, the role of SETD1B in DLBCL remains unknown.

RAB family member 17 (RAB17) is a member of the RAB GTPase family, regulating intracellular vesicle trafficking and membrane dynamics (Sciarra et al. 2024). RAB17 has been implicated in various cellular processes that could contribute to cancer progression (Du et al. 2023). For instance, RAB17 has been shown to regulate the transport of signalling molecules and receptors, potentially influencing cell proliferation, survival, and migration (Wu et al. 2021; Zhou et al. 2021). Given its critical role in intracellular trafficking (Tajbakhsh et al. 2021), RAB17 may regulate the development of DLBCL.

The present study investigated the role SETD1B in DLBCL. The researchers hypothesise that SETD1B-mediated histone modifications may regulate RAB17 expression, thereby affecting intracellular trafficking and contributing to the aggressive phenotype of DLBCL. By elucidating the molecular mechanisms underlying this interaction, the researchers hope to identify potential therapeutic targets for DLBCL.

MATERIAL AND METHODS

Specimen

A total of 32 DLBCL patient samples were collected from Anhui Provincial Children's Hospital between 2020 and 2023. The inclusion criteria were newly diagnosed DLBCL patients with available fresh-frozen tissue samples and complete clinical data. Exclusion criteria included patients with prior treatment for lymphoma, secondary DLBCL, or insufficient tissue for analysis. The study was authorized by the Ethical Committee of Anhui Provincial Children's Hospital. Each patient handed in the informed consent.

Cell Culture

DLBCL cell lines, including SU-DHL-4, OCI-LY10, and DB, were provided by ATCC. Cells were cultured in RPMI-1640 medium containing 10 percent FBS at 37°C in 5 percent CO₂.

Transfection

siRNA targeting RAB17 and SETD1B overexpression plasmids were transfected into OCI-

Y10 cells by Lipofectamine 3000 (Thermo Fisher Scientific).

Quantitative Real-Time PCR (qRT-PCR)

Total RNA was extracted from OCI-Y10 cells. RNA was purified and quantified. cDNA was synthesised by a cDNA Synthesis Kit (Bio-Rad, USA). PCR was conducted by a PCR kit (Bio-Rad, USA). Results were calculated by the $2^{-\Delta\Delta C_t}$ method. GAPDH served as the internal control.

Western Blot

Protein was harvested from OCI-Y10 cells. Protein concentration was calculated. Protein was separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and moved onto polyvinylidene fluoride membranes. After sealing with 5 percent non-fat milk, the membrane was incubated with primary antibodies against SETD1B (ab177288; Abcam, UK), BCL-2 associated X (BAX) (ab32503; Abcam, UK), cl-caspase 3 (cl-CASP3) (ab2302; Abcam, UK), RAB17 (ab155135; Abcam, UK) and α -actin (ab8226; Abcam, UK). Membranes were then incubated with secondary antibodies including goat-anti-rabbit (ab6721; Abcam, UK) and goat-anti-mouse (ab205719). Bands were detected using Western Electrochemiluminescence Blotting Substrate (Bio-Rad).

Chromatin Immunoprecipitation (ChIP) Assay

ChIP was performed to analyse SETD1B-mediated histone modifications, specifically H3K4me3, in DLBCL cell lines. Cells were cross-linked with 1 percent formaldehyde. Chromatin was sonicated to an average fragment size of 200-500 bp. Chromatins were immunoprecipitated using antibodies against H3K4me3 (Abcam, ab8580) and SETD1B (Abcam, ab177288). DNA was extracted from immunoprecipitated chromatin and subjected to qRT-PCR.

Cell Counting Kit-8 (CCK-8) Assay

OCI-Y10 cells were seeded in a 96-well plate. After 48 hours, cells were added with 10 μ L of CCK-8 solution. Optic values were detected by a microplate reader (BioTek).

Flow Cytometry

After transfection, OCI-Y10 cells were stained with Annexin V-FITC and propidium iodide. Apoptotic cells were calculated using a flow cytometry.

Statistical Analysis

Data was analysed using GraphPad8.0 and presented as mean $\pm$ SD. Comparisons were analysed using Student's t-test or one-way ANOVA. $P < 0.05$ were considered statistically significant. All experiments were conducted in triplicate.

RESULTS

SETD1B is Downregulated in DLBCL Patients

SETD1B exerts anti-tumour function in various cancers (Cheon et al. 2022; Friedman et al. 2024). SETD1B was markedly downregulated in DLBCL patients. The area under the ROC curve is 0.891, with 95 percent confidence interval (CI): 0.813~0.969 (Fig. 1A). Therefore, SETD1B mRNA expression in DLBCL patients can be a sensitive biomarker (Fig. 1B). Additionally, downregulated DLBCL predicted advanced stages (Fig. 1C) and distance metastasis (Fig. 1D).




Fig. 1. SETD1B is overexpressed in DLBCL patients

(A) SETD1B mRNA levels in DLBCL patients were analysed by RT-qPCR

(B) ROC curve analysis of A

(C-D) SETD1B mRNA levels in DLBCL patients were analysed by RT-qPCR

** $P < 0.01$, *** $P < 0.001$

SETD1B Drives DLBCL Cell Apoptosis

SETD1B was markedly downregulated in DLBCL cells, particularly in OCI-Y10 cells (Fig. 2A). Therefore, OCI-Y10 cells were used in further study. SETD1B was remarkably upregulated by its overexpression plasmids (Fig. 2B). Overexpressed SETD1B significantly inhibited OCI-Y10 cell viability (Fig. 2C). Overexpressed SETD1B markedly promoted the apoptosis of

OCI-Y10 cells (Fig. 2D) and increased the percentages of TUNEL positive cells (Fig. 2E). Additionally, overexpressed SETD1B remarkably upregulated BAX and cl-CASP3 (Fig. 2F). These results suggested that SETD1B promotes the apoptosis of DLBCL.

SETD1B Upregulates RAB17 Via Promoting Histone Methylation

GSE273152 was used to analysed the differentially expressed genes (DEGs) after SETD1B knockout. RAB17 expression was markedly downregulated (Fig. 3A). Overexpressed SETD1B markedly upregulated RAB17 (Fig. 3B). SETD1B regulates gene expression via mediating histone methylation in its promoter (Hanna et al. 2022; Lin et al. 2025). SETD1B overexpression markedly upregulated the protein expression of H3K4me3 and RAB17 (Fig. 3C). Moreover, the occupancy of H3K4me3 was markedly increased on the promoter of RAB17 (Fig. 3D). However, SETD1B markedly reduced the occupancy of H3K4me3 on the promoter of RAB17 (Fig. 3E). These results suggested that SETD1B regulates RAB17 expression via mediating histone methylation on its promoter in DLBCL.

RAB17 Knockdown Inhibits the Apoptosis of OCI-Y10 Cells

RAB17 expression was markedly reduced by its siRNA (Fig. 4A). RAB17 knockdown significantly promoted OCI-Y10 cell viability (Fig. 4B). Moreover, RAB17 knockdown significantly inhibited OCI-Y10 cell apoptosis (Fig. 4C-E). Moreover, RAB17 knockdown markedly downregulated BAX and cl-CASP3 (Fig. 4F).

DISCUSSION

In the present study, SETD1B was downregulated in DLBCL, which predicted poor clinical outcomes. Interestingly, SETD1B overexpression inhibited the malignant behaviours as well as promoted DLBCL cell apoptosis. Moreover, SETD1B promoted the histone methylation and upregulation of RAB17. However, RAB17 was downregulated in DLBCL. RAB17 knockdown contributed to malignant behaviours of DLBCL cells.




Fig. 2. SETD1B promotes the apoptosis of DLBCL cells (A-B) SETD1B levels were detected by RT-qPCR (C) Cell proliferation was analysed by CCK-8 (D) Cell apoptosis was analysed by flow cytometry (E) Cell death was analysed by TUNEL staining (F) Protein expression was analysed by Western blot ** $P < 0.01$




Fig. 3. SETD1B upregulates RAB17 via promoting histone methylation
 (A) GSE273152 was used to analyse the DEGs after the SETD1B knockout
 (B) mRNA expression was detected using T-qPCR
 (C) Protein expression was detected using Western blot
 (D-E) H3K4me3 methylation of RAB17 was detected using ChIP assay
 ** $P < 0.01$

SETD1B is dysregulated in tumorigenesis. For example, SETD1B mutation contributes to apoptosis-resistance in B cell lymphoma (Sciarra et al. 2024). However, overexpressed SETD1B predicts unsatisfactory outcomes of patients with hepatocellular carcinoma or colorectal cancer (Chen et al. 2019; Lindner et al. 2020). These findings dictated that SETD1B functions as a tumour-suppressor as well as tumour-promoter, which varies with the types of tumours. In this study, downregulated SETD1B in DLBCL patients predicted poor prognosis. However, overexpressed SETD1B promoted DLBCL cell apoptosis.

hallmark of tumorigenesis (Chen et al. 2020; Sutopo et al. 2023). Recent studies have demonstrated that dysregulated histone methylation contributes to the progression of DLBCL (Huang et al. 2021). For instance, SETD1B deficiency-mediated inhibition of methylation of H3K4 promotes the tumour growth of DLBCL (Portelinha et al. 2024). NCAPD3-mediated H3K9 monomethylation on Sirt1 promotes the proliferation of DLBCL (Lu et al. 2025). In this study, downregulated SETD1B inhibited the methylation of H3K4 and induced the progression of DLBCL. Intriguingly, overexpressed SETD1B drove H3K4 methylation and upregulation of RAB17. RAB17








Fig. 4. RAB17 deficiency suppresses DLBCL cell apoptosis
 (A) Protein expression was analysed by Western blot
 (B) Cell proliferation was analysed by CCK-8
 (C-D) Cell apoptosis was analysed by flow cytometry
 (E) Cell death was analysed by TUNEL staining
 (F) Protein expression was analysed by Western blot
 ** $P < 0.01$









exerts anti-tumour function in various cancers. For instance, low levels of RAB17 predict unfavourable outcomes of kidney renal clear cell carcinoma (Zeng et al. 2023). RAB17 deficiency promotes invasion of NSCLC (Wang et al. 2020). However, overexpressed RAB17 suppresses the aggressiveness of ovarian cancer (Guo et al. 2020). In this study, RAB17 expression was reduced in DLBCL. RAB17 knockdown mediated DLBCL cell survival. Therefore, SETD1B may inhibit the progression of DLBCL via regulating RAB17.

However, several limitations of the study should be acknowledged. While the genetic and epigenetic analyses provided valuable sight on the potential of SETD1B in DLBCL, the functional assays were primarily performed in cell lines. Further validation in *in vivo* models is necessary. Additionally, the downstream signalling pathways regulated by SETD1B in DLBCL cells warrant further investigation to fully elucidate the molecular mechanisms underlying its role in apoptosis.

CONCLUSION

In conclusion, SETD1B exerts an anti-tumour function in DLBCL. Overexpressed SETD1B effectively drives DLBCL cell apoptosis. Moreover, SETD1B upregulates RAB17 expression through driving histone methylation on its promoter. However, RAB17 deficiency contributes to the progression of DLBCL.

RECOMMENDATIONS

SETD1B may be a sensitive biomarker of DLBCL. Targeting SETD1B may be a promising strategy for DLBCL.

DECLARATIONS

COMPETING INTERESTS

The authors declare that they have no conflict of interest.

AVAILABILITY OF DATA AND MATERIAL

The data sets used and analysed in the current study are available on reasonable request from the corresponding authors.

FUNDING

Not applicable.

ACKNOWLEDGEMENTS

Not applicable.

CONFLICT OF INTEREST

Not applicable.

REFERENCES

- Chapuy B, Stewart C, Dunford AJ, Kim J, Kamburov A, Redd RA, Lawrence MS, Roemer MGM, Li AJ, Ziepert M, Staiger AM, Wala JA, Ducar MD, Leshchiner I, Rheinbay E, Taylor-Weiner A, Coughlin CA, Hess JM, Pedamallu CS, Livitz D, Rosebrock D, Rosenberg M, Tracy AA, Horn H, van Hummelen P, Feldman AL, Link BK, Novak AJ, Cerhan JR, Habermann TM, Siebert R, Rosenwald A, Thorner AR, Meyerson ML, Golub TR, Beroukhi R, Wulf GG, Ott G, Rodig SJ, Monti S, Neuberg DS, Loeffler M, Pfreundschuh M, Trümper L, Getz G, Shipp MA 2018. Molecular subtypes of diffuse large B cell lymphoma are associated with distinct pathogenic mechanisms and outcomes. *Nat Med*, 24(5): 679-690. doi: 10.1038/s41591-018-0016-8
- Chen D, Li T, Wang C, Lei G, Wang R, Wang Z, Yu L, Yan J, Zhang P, Wang X, Zhang S, Yang P 2019. High level SETD1B gene expression is associated with unfavorable prognosis in hepatocellular carcinoma. *Mol Med Rep*, 19(3): 1587-1594. doi: 10.3892/mmr.2019.9832
- Chen Y, Jiang L, Liu L, Ruan M, Zhou Z, Yang M, Xia L 2022. Advances and challenges in the investigation of metastasis in diffuse large B-Cell Lymphoma. *Mini Rev Med Chem*, 22(22): 2808-2812. doi: 10.2174/1389557522666220519085134
- Chen Y, Ren B, Yang J, Wang H, Yang G, Xu R, You L, Zhao Y 2020. The role of histone methylation in the development of digestive cancers: a potential direction for cancer management. *Signal Transduct Target Ther*, 5(1): 143. doi: 10.1038/s41392-020-00252-1
- Cheon H, Xing JC, Moosic KB, Ung J, Chan VW, Chung DS, Toro MF, Elghawry O, Wang JS, Hamele CE, Hardison RC, Olson TL, Tan SF, Feith DJ, Ratan A, Loughran TP 2022. Genomic landscape of TCR α and TCR γ T-large granular lymphocyte leukemia. *Blood*, 139(20): 3058-3072. doi: 10.1182/blood.2021013164
- Dabrowska-Iwanicka A, Nowakowski GS 2024. DLBCL: Who is high risk and how should treatment be optimized? *Blood*, 144(25): 2573-2582. doi: 10.1182/blood.2023020779
- Du H, Li S, Lu J, Tang L, Jiang X, He X, Liang J, Liao X, Cui T, Huang Y, Liu H 2023. Single-cell RNA-seq and bulk-seq identify RAB17 as a potential regulator of angiogenesis by human dermal microvascular endothelial cells in diabetic foot ulcers. *Burns Trauma*, 11: tkad020. doi: 10.1093/burnst/tkad020

- Friedman CF, Manning-Geist BL, Zhou Q, Soumerai T, Holland A, Da Cruz Paula A, Green H, Ozsoy MA, Iasonos A, Hollmann T, Leitao MM Jr, Mueller JJ, Makker V, Tew WP, O'Cearbhaill RE, Liu YL, Rubinstein MM, Troso-Sandoval T, Lichtman SM, Schram A, Kyi C, Grisham RN, Causa Andrieu P, Wherry EJ, Aghajanian C, Weigelt B, Hensley ML, Zamarin D 2024. Nivolumab for mismatch-repair-deficient or hypermutated gynecologic cancers: A phase 2 trial with biomarker analyses. *Nat Med*, 30(5): 1330-1338. doi: 10.1038/s41591-024-02942-7
- Guo M, Li S, Zhao X, Yuan Y, Zhang B, Guan Y 2020. Knockdown of circular RNA Hsa_circ_0000714 can regulate RAB17 by sponging miR-370-3p to reduce paclitaxel resistance of ovarian cancer through CDK6/RB Pathway. *Oncotargets Ther*, 13: 13211-24. doi: 10.2147/OTT.S285153
- Hanna CW, Huang J, Belton C, Reinhardt S, Dahl A, Andrews S, Stewart AF, Kranz A, Kelsey G 2022. Loss of histone methyltransferase SETD1B in oogenesis results in the redistribution of genomic histone 3 lysine 4 trimethylation. *Nucleic Acids Res*, 50(4): 1993-2004. doi: 10.1093/nar/gkac051
- Huang YH, Cai K, Xu PP, Wang L, Huang CX, Fang Y, Cheng S, Sun XJ, Liu F, Huang JY, Ji MM, Zhao WL 2021. CREBBP/EP300 mutations promoted tumor progression in diffuse large B-cell lymphoma through altering tumor-associated macrophage polarization via FBXW7-NOTCH-CCL2/CSF1 axis. *Signal Transduct Target Ther*, 6(1): 10. doi: 10.1038/s41392-020-00437-8
- Kotlov N, Bagaev A, Revuelta MV, Phillip JM, Cacciapuoti MT, Antysheva Z, Svekolkin V, Tikhonova E, Mihecheva N, Kuzkina N, Nos G, Tabbo F, Frenkel F, Ghione P, Tsiper M, Almog N, Fowler N, Melnick AM, Leonard JP, Inghirami G, Cerchetti L 2021. Clinical and biological subtypes of B-cell lymphoma revealed by microenvironmental signatures. *Cancer Discov*, 11(6): 1468-1489. doi: 10.1158/2159-8290.CD-20-0839
- Kranz A, Anastassiadis K 2020. The role of SETD1A and SETD1B in development and disease. *Biochim Biophys Acta Gene Regul Mech*, 1863(8): 194578. doi: 10.1016/j.bbagr.2020.194578. PubMed PMID: 32389824.
- Lin Z, Rong B, Lyu R, Zheng Y, Chen Y, Yan J, Wu M, Gao X, Tang F, Lan F, Tong MH 2025. SETD1B-mediated broad H3K4me3 controls proper temporal patterns of gene expression critical for spermatid development. *Cell Res*, 35(5): 345-361. doi: 10.1038/s41422-025-01080-0
- Lindner P, Paul S, Eckstein M, Hampel C, Muenzner JK, Erlenbach-Wuensch K, Ahmed HP, Mahadevan V, Brabletz T, Hartmann A, Vera J, Schneider-Stock R 2020. EMT transcription factor ZEB1 alters the epigenetic landscape of colorectal cancer cells. *Cell Death Dis*, 11(2): 147. doi: 10.1038/s41419-020-2340-4
- Lu T, Yang J, Cai Y, Ding M, Yu Z, Fang X, Zhou X, Wang X 2025. NCAPD3 promotes diffuse large B-cell lymphoma progression through modulating SIRT1 expression in an H3K9 monomethylation-dependent manner. *J Adv Res*, 68: 163-178. doi: 10.1016/j.jare.2024.02.024
- Nakhoda S, Vistarop A, Wang YL 2023. Resistance to Bruton tyrosine kinase inhibition in chronic lymphocytic leukaemia and non-Hodgkin lymphoma. *Br J Haematol*, 200(2): 137-149. doi: 10.1111/bjh.18418
- Ozyerli-Goknar E, Kala EY, Aksu AC, Bulut I, Cingöz A, Nizamuddin S, Biniossek M, Seker-Polat F, Morova T, Aztekin C, Kung SHY, Syed H, Tuncbag N, Gönen M, Philpott M, Cribbs AP, Acilan C, Lack NA, Onder TT, Timmers HTM, Bagci-Onder T 2023. Epigenetic-focused CRISPR/Cas9 screen identifies (absent, small, or homeotic)2-like protein (ASH2L) as a regulator of glioblastoma cell survival. *Cell Commun Signal*, 21(1): 328. doi: 10.1186/s12964-023-01335-6
- Pi M, Kuang H, Yue C, Yang Q, Wu A, Li Y, Assaraf YG, Yang DH, Wu S 2022. Targeting metabolism to overcome cancer drug resistance: A promising therapeutic strategy for diffuse large B cell lymphoma. *Drug Resist Updat*, 61: 100822. doi: 10.1016/j.drug.2022.100822
- Portelinha A, Wang S, Parsa S, Jiang M, Gorelick AN, Mohanty S, Sharma S, de Stanchina E, Berishaj M, Zhao C, Heward J, Aryal NK, Tavana O, Wen J, Fitzgibbon J, Dogan A, Younes A, Melnick AM, Wendel HG 2024. SETD1B mutations confer apoptosis resistance and BCL2 independence in B cell lymphoma. *J Exp Med*, 221(10). doi: 10.1084/jem.20231143
- Roston A, Evans D, Gill H, McKinnon M, Isidor B, Cogné B, Mwenifumbo J, van Kamebeek C, An J, Jones SJM, Farrer M, Demos M, Connolly M, Gibson WT; CAUSES Study; EPGEN Study 2021. SETD1B-associated neurodevelopmental disorder. *J Med Genet*, 58(3): 196-204. doi: 10.1136/jmedgenet-2019-106756
- Schmitz R, Wright GW, Huang DW, Johnson CA, Phelan JD, Wang JQ, Roulland S, Kasbekar M, Young RM, Shaffer AL, Hodson DJ, Xiao W, Yu X, Yang Y, Zhao H, Xu W, Liu X, Zhou B, Du W, Chan WC, Jaffe ES, Gascoyne RD, Connors JM, Campo E, Lopez-Guillermo A, Rosenwald A, Ott G, Delabie J, Rimsza LM, Tay Kuang Wei K, Zelenetz AD, Leonard JP, Bartlett NL, Tran B, Shetty J, Zhao Y, Soppet DR, Pittaluga S, Wilson WH, Staudt LM 2018. Genetics and pathogenesis of diffuse large B-Cell lymphoma. *N Engl J Med*, 378(15): 1396-1407. doi: 10.1056/NEJMoa1801445
- Sciarra R, Merli M, Cristinelli C, Lucioni M, Zibellini S, Riboni R, Furlan D, Uccella S, Zerbi C, Bianchi B, Gotti M, Ferretti VV, Varraso C, Fraticelli S, Lazic T, De-francesco I, Mora B, Libera L, Mazzacane A, Carpi F, Berliner M, Neri G, Rizzo E, De Paoli F, Sessa F, Passamonti F, Paulli M, Arcaini L 2024. Molecular characterization of diffuse large B-cell lymphomas associated with hepatitis C virus infection. *Br J Haematol*, 204(6): 2242-2253. doi: 10.1111/bjh.19378
- Shimada K, Yoshida K, Suzuki Y, Iriyama C, Inoue Y, Sanada M, Kataoka K, Yuge M, Takagi Y, Kusumoto S, Masaki Y, Ito T, Inagaki Y, Okamoto A, Kuwatsuka Y, Nakatochi M, Shimada S, Miyoshi H, Shiraishi Y, Chiba K, Tanaka H, Miyano S, Shiozawa Y, Nannya Y, Okabe A, Kohno K, Atsuta Y, Ohshima K, Nakamura S, Ogawa S, Tomita A, Kiyoi H 2021. Frequent genetic alterations in immune checkpoint-related genes in intravascular large B-cell lymphoma. *Blood*, 137(11): 1491-1502. doi: 10.1182/blood.2020007245
- Sutopo NC, Kim JH, Cho JY 2023. Role of histone methylation in skin cancers: Histone methylation-modifying enzymes as a new class of targets for skin cancer treatment. *Biochim Biophys Acta Rev Cancer*, 1878(3): 188865. doi: 10.1016/j.bbcan.2023.188865

- Tajbakhsh A, Farahani N, Gheibihayat SM, Mirkhabbaz AM, Savardashtaki A, Hamblin MR, Mirzaei H 2021. Autoantigen-specific immune tolerance in pathological and physiological cell death: Nanotechnology comes into view. *Int Immunopharmacol*, 90: 107177. doi: 10.1016/j.intimp.2020.107177
- Wang M, Wang W, Ding J, Wang J, Zhang J 2020. Downregulation of Rab17 promotes cell proliferation and invasion in non-small cell lung cancer through STAT3/HIF-1 α /VEGF signaling. *Thorac Cancer*, 11(2): 379-388. doi: 10.1111/1759-7714.13278
- Wu S, Luo M, To KKW, Zhang J, Su C, Zhang H, An S, Wang F, Chen D, Fu L 2021. Intercellular transfer of exosomal wild type EGFR triggers osimertinib resistance in non-small cell lung cancer. *Mol Cancer*, 20(1): 17. doi: 10.1186/s12943-021-01307-9
- Zeng Z, Zhang Z, Cheng X, Yang H, Gong B, Zhou X, Zhang C, Wang X, Wang G 2023. Downregulation of RAB17 have a poor prognosis in kidney renal clear cell carcinoma and its expression correlates with DNA methylation and immune infiltration. *Cell Signal*, 109: 110743. doi: 10.1016/j.cellsig.2023.110743
- Zhou X, Xia G, Liu Y et al. 2021. Analysis of carcinogenic signaling networks in endometrial cancer identifies RAB17 as a potential target. *J Cell Physiol*, 236(1): 328-339. doi: 10.1002/jcp.29845

Paper received for publication in June, 2025
Paper accepted for publication in January, 2026