

Inherently Delayed Germination of Elgon Teak (*Olea capensis*) Seeds in Kakamega Forest, Kenya

Mugatsia Tsingalia¹ and Nyongesa Humphrey²

¹*Department of Biological Sciences, Masinde Muliro University of Science and Technology,
P.O. Box Kakamega Kenya*

E-mail: mugatsi2005@yahoo.com

²*Institute of Sugar Technology, Masinde Muliro University of Science and Technology*

P.O. Box 190 – 50100, Kakamega, Kenya

E-mail: hnyongesa2005@yahoo.com

KEYWORDS Bioassay. Seedlings. Seed Germination. Elgon Teak. Kakamega Forest. Allelopathy. Autotoxicity

ABSTRACT *Olea capensis* in Kakamega forest does not show any evidence of natural regeneration under the parent trees. This study sought to determine the factors that may be responsible for the lack of natural regeneration under the adult parent crowns in the Kakamega forest, Kenya. The study was carried out between November, 1999 and August, 2000 which marked the end of the fruiting period when one would expect to find a lot of germinated seeds on the forest floor. Contrary to this expectation, there was no observed germination. It was, therefore, envisaged that chemical inhibition by the parent tree may be responsible for the lack of seed germination and establishment under the parent crowns. Three principal bioassay methods were used to test whether chemical interaction indeed may be a possible mechanism inhibiting the germination and establishment of seeds and seedlings in the immediate vicinity of *Olea* adults inside the forest. To distinguish allelopathic interactions from others such as seed inviability, bioassays were applied on experimentally germinated seeds in the nearby forest tree nursery where sprouting of buds and shoots was monitored and measured. Controls were watered with distilled water. Live-shoot bioassay significantly inhibited seed germination. Both shoot and root leachates significantly ($p \leq 0.05$) reduced shoot and root budding in young experimental seedlings in the forest tree nursery. Shoot leachates appear to be the most effective inhibitors of seed germination and establishment of *Olea* seeds and seedlings inside the Kakamega forest. Shoot leachates also significantly ($p \leq 0.05$) retarded the growth of *Olea* seedlings. This has practical implications for the regeneration of *Olea* in the Kakamega forest and explains the clumped distribution of *Olea* adults in the forest. Inhibitory substances appear to delay germination of seeds for periods long enough for predation to occur. The implication is, only seeds that are dispersed distances away from *Olea* adults will germinate and establish, and the dispersal pattern determines the eventual spatial distribution.