

## **Genetic Studies on Eleven Populations of Jammu and Kashmir, India**

**H. Walter<sup>1</sup>, M.K. Bhasin<sup>2</sup>, Heldi Danker-Hopfe<sup>1</sup>, Asha Khanna<sup>2</sup>, Ute Hahm<sup>1</sup> and  
Iris Wittl<sup>1</sup>**

*<sup>1</sup>Department of Human Biology, University of Bremen, D-2800 Bremen 33, Germany*

*<sup>2</sup>Department of Anthropology, University of Delhi, Delhi 110 007, India*

**KEYWORDS** Genetic Markers. Genetic Variability and Distances. Jammu and Kashmir

**ABSTRACT** Eleven population groups of Jammu and Kashmir (India) with a total of n = 1131 unrelated male and female individuals were typed for HP polymorphism and for GC and PI subtype polymorphisms. The genetic variability among these eleven population groups under study is considerable, which is analyzed in detail. One of the most striking observations is the distribution of PI allele frequencies, in which the three Tibeto-Chinese speaking population groups – Baltis, Bodhs and Tibetans – show remarkable differences from all the other groups under study. Genetic distance analyses revealed a genetic distance pattern, which is in good accordance with the ethnic structure and ethnohistory of the populations of this region.