

## **Chemical Evaluation of South African Prickly Pear Cladodes Varieties as Fodder for Ruminants Grown in Mara ADC, South Africa**

**Mokoboki Kwena<sup>1,2\*</sup>, Sebola Nthabiseng<sup>1,2</sup> and Makgobatlou Grace<sup>3</sup>**

*<sup>1</sup>Animal Science Programme, North West University, Mafikeng Campus,  
Mmabatho 2735, Private Bag x 2046, South Africa*

*<sup>2</sup>Food Security and Safety Niche Area, School of Agricultural Sciences, Faculty of Agriculture,  
Science and Technology, North-West University, Mmabatho, Mafikeng 2735, South Africa*

*<sup>3</sup>Mara Research Station, Private Bag x 2467, Makhado, 0920, South Africa  
Mobile: \*+27769562801>, E-mail: \*<21576130@nwu.ac.za>*

**KEYWORDS** Cladodes. Fiber. Protein. Chemical Evaluation. Prickly Pears

**ABSTRACT** The aim of the study was to determine the chemical composition of South African prickly pears cladodes varieties as fodder for ruminants in Mara ADC, South Africa. The experimental field was divided into four replications with ten different prickly pears varieties within each replication. The following results were obtained (percent of dry matter): crude protein 6.23-10.63, neutral detergent fiber 19.36-38.29, acid detergent fiber 12.25-19.25, ether extract 1.19-2.62 and the phosphorus content 0.19-0.29. It was concluded that the environment aspects in the Limpopo Province did not negatively impact the chemical composition of prickly pears since the concentrations of fiber was lower and crude protein content was higher.