

## **Vitamin D Dependent Rickets Type II with Alopecia**

**M. Vasudeva Murali and T. Usharani**

*Gandhi Hospital, Department of Pediatrics), Musheerabad, Secunderabad 500 003,  
Andhra Pradesh, India*

**KEYWORDS** VDDR II. Rickets. Hair Loss. 1,25-dihydroxyvitamin D<sub>3</sub>. Vitamin D Receptor

**ABSTRACT** Vitamin D dependent rickets Type II (VDDR II) is a rare form of autosomal recessive disorder, which is inherited and often caused by mutation of a gene on vitamin D receptor which in turn prevents normal physiological response to 1, 25(OH)<sub>2</sub> vitamin D. The elevated circulating levels of 1,25-dihydroxyvitamin D<sub>3</sub> is a diagnostic feature of VDDR II, while the presence of alopecia differentiates it from vitamin D-dependent rickets Type I. VDDR II is considered to be the best example of aberrant target tissue response. The researchers thereby report a case of a two-year old female child with short stature, typical features of rickets and lack of hair growth. Upon investigation the level of 1,25(OH)<sub>2</sub>D was found to be high which is diagnostic of VDDR type II. The child was started on high dose of calcitriol and oral calcium but on follow-up did not show much improvement probably due to poor compliance.