

Noise Pollution and Human Health in Trabzon Parks

Buket Ozdemir¹, Elif Bayramoglu² and Oner Demirel³

*Forestry Faculty, Department of Landscape Architecture, Karadeniz Technical University,
61080 Trabzon, Turkey*

E-mail: ¹<buketozdemir@hotmail.com>, ²<elifsol_@hotmail.com>, ³<odofe01@yahoo.com>

KEYWORDS Traffic Noise. Plant Barriers. City Parks. Public Health

ABSTRACT Since the 1980s, rapid and unplanned urbanization has caused environmental pollution. It is accepted that noise pollution has a major impact on health, such as physical, physiological, psychological and performance-related effects, all over the world. This paper provides an evaluation of noise pollution in three urban parks in the city of Trabzon, located in the north-eastern part of Turkey. Equivalent noise point levels were measured during a 3-minute spread at each park. Measured values were compared with national legislation (Law: 60) allowed limits, and the parks were thus classified as either "acoustically polluted or unpolluted." Urban parks in Trabzon's city center, surrounded by roads with heavy traffic and intense commercial activities, do not satisfy any of the standards used. The most noise-polluted parks in Trabzon were the Meydan Park, Atapark and Fatih Park with measured Leq of dB (A) Meydan park (63.74), dB (A) Atapark (64.15) and dB (A) Fatih park (64.67). This paper examines the opportunity of using plant material (*Syringa vulgaris*, *Viburnum lantana* and *Acer pseudoplatanus*) to minimize noise to acceptable levels in the three urban parks located inside Trabzon's city center. At the end of this examination, it was concluded that is possible to reduce noise to acceptable levels in all three parks, by using green barriers decorated with plant material. In this way, the noise level was reduced, which will have a positive effect on human health.

INTRODUCTION

Environmental pollution becomes more severe and widespread due to population growth, urbanization and industrialization in the cities (Ralte et al. 2013). There are many factors which cause the environment to be polluted and one of those undesired and unpleasant factors is 'noise' which affects the quality of life (Haq et al. 2014). Numerous researchers have demonstrated that exposure to environmental noise may increase the risks related to personal health, such as nervous frailty, extreme irritability, muscle cramps, stress and anxiety, dizziness, headache and migraine, anger, loss of body balance. Consequently, noise pollution is one of the major problems for developing countries. There is a need to control the noise exposure levels in sensitive areas such as hospitals, schools and kindergartens (Mitra 2008; Oyedepo and Abdullahi

2009; Noori and Zand 2013; Amin et al. 2014; Marriscal-Rammires et al. 2014; Mukhola 2014).

This situation has led to protective, noise-controlling legislation, such as the Occupational Safety and Health Act, the Noise Control Act, European Directive 2002/49/EC and the Quiet Communities Act (Moudon 2009). The World Health Organization (WHO) and the US Environmental Protection Agency consider that the safe level of equivalent continuous sound (LAeq) for human health is 70 decibels (dB) (WHO 2000). However, according to this legislation, 80 million people suffer from unacceptable noise levels and 170 million experience serious annoyances during daytime in the European Union (EU) (Miedema 2007).

The urban parks, which are the main focus of this paper, stand out as important areas of social life providing access to nature and opportunities for various uses (rest, sports, relaxation, games, cultural events and sightseeing) with acoustic comfort, with a large number of studies focusing on the problem of noise pollution at urban parks (Li et al. 2002; Morillas et al. 2002; Zannin and Szermetta 2003; Lam et al. 2005; Cengiz et al. 2012; Hunashal and Patil 2012).

In addition to this there are many field surveys about the reduction of noise using plants (Beckett et al. 2000; Fang and Ling 2005; Ozer et

Address for correspondence:

Dr. Elif Bayramoglu
Forestry Faculty,
Department of Landscape Architecture,
Karadeniz Technical University,
61080 Trabzon, Turkey
Telephone: +90 462 377 4083,
Fax: +90 462 325 7499
E-mail: elifsol_@hotmail.com

al. 2008; Pathak et al. 2008; Islam et al. 2012; Vasilakopoulou et al. 2014).

In Turkey, noise is now considered as one of the main environmental problems (Kelkit 2003). The number of cars and levels of urban noise in Turkey have been increasing and reached high levels (Bekci et al. 2013). Due to the increase in population density and the process of development as a city, Trabzon cannot provide the acoustic comfort of living to its public and continues to present unhealthy living conditions. Because of insufficient communication and the huge number of cars in traffic, the noise problem has become more difficult to solve. This paper was carried out in order to determine the extent of noise levels in Trabzon's urban parks (Meydan Park, Fatih Park, Atapark). The second aim of the paper was to explore solving this problem with the use of plants.

Noise and Health

Today, noise pollution is generally defined as unwanted or loud noise or undesirable sound levels. Environmental noise may cause adverse effects for individuals as well as creating situational factors. Noise pollution is distinguished from other pollution categories due to its source and diffusion characteristics, which can adversely affect public health and environmental quality in the urban environment. The WHO recently estimated that traffic noise could conservatively account for over one million health years of life lost annually in the European Union and Western European countries. At the same time WHO recognized the following effects on the health of the population that can emanate from noise: disturbance in sleep patterns, cardio respiratory and psycho-physiological systems and

hearing. It may also have negative and intervening effects on communication, productivity and social behavior (WHO 1993, 2011; Tsitsoni et al. 2005; Samara and Tsitsoni 2007).

Noise can be defined as the level of sound, which exceeds the acceptable level and creates annoyance. Frequent exposure to high level of noise can cause severe stress on the auditory and nervous system (Subramani et al. 2012). The impact of noise on human health has been studied in four aspects. These are as follows: physical effects (temporary or permanent hearing loss), physiological effects (breathing difficulties, heart disorders, high blood pressure, sleep disturbance), psychological effects (adverse emotions including anger, anxiety, depression and behavioral disorders) and performance-related effects (reduction in reading, learning and work performance, lack of concentration) (Onder and Kocbeker 2012). The effect of noise on children was also reported including findings such as increased blood pressure and annoyance reactions (Paunovic et al. 2009; Liu et al. 2014). Some studies conducted on children describe effects of noise on executive functioning (EF) (decision making, working memory and self-regulation of emotions and behaviors) and cognitive performance (Fritschi et al. 2011; Belojevic et al. 2012).

The effect of noise is the most prominent feature of the noise from the physical of inability to hear. Most studies recommend that for L_{Aeq} , exposure to 24h of less than 70dB does not lead to any permanent hearing loss. At the same time 70dB (L_{Aeq} , 24h) noise conditions can cause permanent damage to human health (King and Davis 2003). According to the Ministry of Environment and Forestry standards found in the regulation (2002/49/EC), the levels of compliance are given in Table 1.

Table 1: Standards of noise level for various areas of community (Environmental Noise Assessment Regulation)

Description of area	Noise level dB(A)			
	Refurbished ways		Existing roads	
	Day time 6.00AM-9.00PM	Night time 9.00PM-6.00AM	Day time 6.00AM-9.00PM	Night time 9.00PM-6.00AM
Rural area	55	45	60	50
Sensitive Areas (parks, schools, hospitals, mosques, Silence area)	60	50	65	55
Settled Area	63	53	68	58
Commercial Area	65	55	75	75
Industrial Area	75	70	75	75

Noise and Plants

Since plants, which are living organisms, can reduce noise, they are used in landscape architecture as “natural, living noise barriers.” Accordingly, green natural elements found in the cities are required for health reasons as they make positive visual and psychological contributions to human health and well-being (Ozbilen and Var 1992; Bekci et al. 2012). The initial studies on the noise prevention using plants were conducted by Beck and Meyer in 1960-1970s. Some types of plants are better at performing noise reduction than others. Bernatzky (1978) state that pollution-preventing characteristic of plant materials change depending on the plant variety, that is, needle-leaf or evergreen plant, its crown diameter and the size and hardness of its leaves. Especially tree belts along roads can be used as solutions to achieve road traffic noise reduction but few quantitative data have been reported on the significance of height, density, width and length of tree belts for noise reduction (Fang and Ling 2003, 2005; Renterghem 2014). Despite the limited knowledge available plants have been used as plant barriers while trees have been used as tree belts in the landscape architecture for the purpose of noise reduction.

MATERIAL AND METHODS

Material

The study area was the city of Trabzon located on the Black Sea coast in the north-east of Turkey (N 41°00'00" and E 39°43'00"). The city of Trabzon's surface area is 4.865 km² and total population is 757.898. People living in the urban, the centres of districts, comprises 426.882 of this population. As the majority of the population lives in city centres, urban parks are used extensively. This paper was conducted in three urban parks in the city of Trabzon which are presented in Figure 1. The parks, Meydan Park, Fatih Park and Atapark, are located along the Tanjant Highway which is prone to traffic jam problems.

As given in Table 2, Meydan Park is located in the center of the city. It is a strictly urban park, surrounded by roads with heavy traffic. Fatih Park, which is 200 metres away from the city center, is the only park with a children's playground and is also surrounded by roads. Atapark is in 4 km of walking distance to the city center. The most important feature of this park is that it is surrounded by different architectural works such as the ancient Trabzon Castle and the city walls, historical Gülbahar Hatun Muse-




Fig. 1. Study areas

Table 2: Description of study area

Name	Location	Characteristics	Plantation	Facilities
<i>Fatih Park</i>	Distance 200m to the city center	4764 m ² hard surface 5016 m ² green areas 9780 m ² total area 350 m ² playground 155 sitting units	128 tall trees 20 medium trees 410 small trees 132 m ² seasonal flower area	Police station Playground Wetland areas
<i>Meydan Park</i>	In the city center.	4270 m ² hard surface 1500 m ² green areas 5770 m ² total area 184 sitting units	74 tall trees 25 medium trees 495 small trees 210 m ² seasonal flower area	Mosque City hall Shopping center Ceremony area Wetland areas
<i>Atapark</i>	Neighbor to Gülbaharhatun. Distance 4 km to the city.	4939 m ² hard surface 2115 m ² green areas 4939 m ² areas used by business operator on hard surface 713 m ² playground 104 sitting units	147 tall trees 7 medium trees 341 small trees 72m ² seasonal flower 7767 m ² total area	Mosque Library Hospital Shopping centre Theatre Wetland areas

um, Varliba° Shopping Mall and Trabzon Central Library. Additionally, it is subject to heavy traffic.

Methods

To detect the extent to which noise can be reduced with the help of plant elements in the urban parks with heavy traffic, measurements were made at 31 points in total in three urban parks in Trabzon (10 points in Meydan Park, 13 points in Atapark, 8 points in Fatih Park). Measurement points were decided considering the 8:00 am – 5:00 pm time interval, the working time, when traffic was heavy. Measurements were made on Mondays, the first working day of the

week. Due to the proximity of the parks to the roads of heavy traffic, measurements were made at 3 minute period intervals at the corner and central points of the parks surrounded by roads. Measurements were performed at the height of 1.20 m, taking as a measurement basis the average ear height of a person in seated position. Plants chosen to prevent noise in the scope of the paper were preferred due to their capacity to absorb noise as a result of physical characteristics such as dense foliage and branching, plants with leaves or coniferous plants (adapted to the climate conditions of Trabzon), plants with strong stem structure and those that require minimal ecological conditions. In this scope, *Syringa vulgaris* (Class IV), *Viburnum*

Table 3: Suitable to the climatic conditions of Trabzon plant species

Latin name	Class	Crown diameter (mt)	Height (mt)
<i>Acer negundo</i>	Class III(4-6 dB)	6-8	10-25
<i>Betula pendula</i>		8-10	25-30
<i>Lonicera japonica</i>	Class IV(6-8 dB)	dispersed	2-3
<i>Forsythia intermedia</i>		dispersed	1.5-2
<i>Cornus alba</i>		3-4	4-5
<i>Syringa vulgaris</i>	Class V(8-10 dB)	4	6-7
<i>Ilex aquifolium</i>		3	8
<i>Quercus robur</i>		15-20	50
<i>Rhododendron ponticum</i>		dispersed	5-6
<i>Viburnum lantana</i>	Class VI(10-12 dB)	3	3-4
<i>Viburnum rhytidophyllum</i>		3	3
<i>Tilia platyphyllos</i>		10-12	20-30
<i>Acer pseudoplatanus</i>		25	20-30

lantana (Class V), and *Acer pseudoplatanus* (Class VI) plants listed in Table 3 were selected as study plants, since they could adapt to regional climate conditions to be used as plant materials in the mentioned parks with the aim to reduce noise level of studied parks down to normal dB levels. As in Beck’s (1967) study, noise barriers were organized as noise-preventing green barriers in 3-component plant groups of “tree-small tree-bush,” increasing at vertical axis from small to large component. or Meydan Park, $L_{Aeq}=64.67$ for Fatih Park and $L_{Aeq}=64.15$ for Atapark. As a result of these findings, all studied urban parks were concluded to have exceeded the noise level. In order to reduce the noise down to acceptable levels and to keep these areas within the limits of Noise Control Regulations, the noise level was reduced using plant elements similar to those used by Fang and Ling (2005), Tyagi et al. (2006), Yang et al. (2011), Al-Dabbous and Kumar (2014).

Three-plant groups, which are given in Figure 2, *Syringa vulgaris* (Sv), *Viburnum lantana* (Vl), and *Acer pseudoplatanus* (Ap) were arranged in this study similar to that in Fang and Ling (2003) and Beck (1967) studies. In studies where similar plant groups were used Fang and Ling (2003) showed the noise levels to be reduced by 2.9-6 dB(A), Renterghem et al. (2012) 3-4.7 dB(A) and Renterghem et al. (2014) by 1.1-3.6 dB(A). In this study as in Huddart (1990),

RESULTS AND DISCUSSION

Noise level measurements made in the urban parks of Trabzon city were evaluated for each park separately. Evaluation results are listed in Table 4. Measurement points were associated with the proximity of parks to roads and their surface area; thus, more measurements were made in the areas with heavy traffic.

Measurements produced the following results for the studied urban parks: $L_{Aeq}=63.74$ these plant groups were recorded to reduce noise by 9 dB(A). This value was calculated by averaging the noise reduction values of the studied plant groups (listed in Table 3) by using the following formula:

$$Sv((6+8)/2) + Vl((8+10)/2) + Ap((10+12)/2) = 7+9+11 = 27/3 = 9 \text{ dB(A)}$$

New dB(A) values were calculated for each point separately on the basis of the selected plant groups. Resulting values are presented in Table 5.

According to Table 5, the noise level of Meydan Park was reduced to $L_{Aeq}=54.74$ while that of Fatih Park was lowered to $L_{Aeq}=55.67$ and for Atapark to $L_{Aeq}=55.15$ dB(A). These values were found to be close to the values obtained from the studies measuring noise level (Ahmad et al. 2006; Zannin et al. 2006; Ozer et al. 2008) and were approximated to the limits stipulated by the Noise Control Regulation.

Table 4: Sound levels measured in each park (between 9:00 am – 10:30 am)

Measurement points	Urban parks								
	Meydan Park			Fatih Park			Atapark		
	L_{AMax}	L_{AMin}	L_{Aeq}	L_{AMax}	L_{AMin}	L_{Aeq}	L_{AMax}	L_{AMin}	L_{Aeq}
1	70.1	56.2	60.7	72.3	61.8	65.4	85	65.1	71.8
2	70	56	62.6	72.2	59.6	63.6	71.7	61.1	67.1
3	70	59.9	63.7	85	59	72.5	68.8	58.9	63.9
4	73.2	63.5	66.7	66.8	55.5	60.2	74.5	60.3	67.4
5	72.7	56	64.4	73.9	55.3	65.1	70.3	57.2	63.2
6	73.7	60.5	63.8	69.2	53.5	58.4	71.1	54.9	60.9
7	69.5	58.5	64.1	71.7	54.3	62.3	70.9	50.6	58.3
8	70.8	59.4	65.8	89.6	56.6	69.9	72.9	53.7	63.8
9	72.9	58	61.6				65	53.6	57.3
10	77.2	61.4	64				67.1	56.1	61.4
11							71.9	55.7	64.7
12							80.3	62.3	69.2
13							71	57.9	65

L_{AMax} : maximum sound level, L_{Aeq} : equivalent continuous sound level, L_{AMin} : minimum sound level




Fig. 2. The sample of group plants for the city of Trabzon

CONCLUSION

Traffic density increases in parallel with the global population growth, which affects noise pollution in urban parks. In this scope, it is possible to reduce the noise levels of park areas to

normal levels by using natural or artificial landscape elements. Individual or in-group use of plant elements can reduce noise levels. Considering use density of urban parks, reduction of the noise levels of these parks down to normal noise levels will offer positive contributions

Table 5: New noise level decreases with 9dB(A)

Measurement points	Urban parks								
	Meydan Park			Fatih Park			Atapark		
	L_{AMax}	L_{AMin}	L_{Aeq}	L_{AMax}	L_{AMin}	L_{Aeq}	L_{AMax}	L_{AMin}	L_{Aeq}
1	61.1	47.2	51.7	63.3	52.8	56.4	76	56.1	62.8
2	61	47	53.6	63.2	60.6	54.6	62.7	52.1	58.1
3	61	50.9	54.7	76	60	63.5	59.8	49.9	54.9
4	64.2	54.5	57.7	57.8	46.5	51.2	65.5	51.3	58.4
5	63.7	47	55.4	64.9	46.3	56.1	61.3	48.2	54.2
6	64.7	51.5	54.8	60.2	44.5	49.4	62.1	45.9	51.9
7	60.5	49.5	55.1	62.7	45.3	53.3	61.9	41.6	49.3
8	61.8	50.4	56.8	80.6	47.6	60.9	63.9	44.7	54.8
9	63.9	49	52.6				56	44.6	48.3
10	68.2	52.4	55				58.1	47.1	52.4
11							62.9	46.7	55.7
12							71.3	53.3	60.2
13							62	48.9	56

L_{AMax} : maximum sound level, L_{Aeq} : equivalent continuous sound level, L_{AMin} : minimum sound level

for users in physical, physiological and performance-related terms.

The use of plant materials in especially in sensitive areas will serve as precautions for potential public health problems, specifically those concerning children.

Although noise pollution is the most frequent type of pollution in Turkey, it is also the less cared for as well as dealt least with using legal action. Even though the legal framework is suitable, the issue is not given much importance. This result reveals the necessity for the cooperation of the governmental institutions and non-governmental organizations for efforts against noise and for reduction of negative effects of noise levels.

RECOMMENDATIONS

Related public agencies and institutions should develop projects to raise public awareness for noise reduction. Since it is impossible to control the vehicle density, which increases in parallel with the population growth, the practice to reduce noise by using living plant barriers should be generalized, particularly to the urban parks, children's playgrounds and open space areas. In order to obtain the anticipated results from the use of these plant species, care should be taken with the properties of the plant material and their application principles. In order to reduce noise, increasing the distance between living areas (hospitals, parks, schools, etc.) and the heavy-traffic roads and use of plant barriers should be the priority.

REFERENCES

- Ahmad J, AL-Omarý A, Sharabi R 2006. Evaluation of traffic noise pollution in Amman, Jordan. *Environmental Monitoring and Assessment*, 120: 499-525.
- Alam JB, Alam MJB, Rahman MM, Dikshit AK, Khan SK 2006. Study on traffic noise level of Sylhet by multiple regression analysis associated with health hazards. *Iran Journal of Environment Health Science Engineering*, 3: 71-78.
- Al-Dabbous AN, Kumar P 2014. The influence of roadside vegetation barriers on airborne nanoparticles and pedestrians exposure under varying wind conditions. *Atmospheric Environment*, 90: 113-124.
- Amin N, Sikder I, Zafor NA, Chowdhury MAI 2014. Assessment of noise pollution of two vulnerable sites of Sylhet City, Bangladesh. *International Journal of Water Resources and Environmental Engineering*, 6: 112-120.
- Beck G 1967. *Pflanzen als mittel zur laermbekaempfung*. Berlin: Patzer – Verlag.
- Beckett KP, Freer-Smith PH, Taylor G 2000. Particular pollution capture urban trees: Effect of species and wind speed. *Global Change Biology*, 6: 995-1003.
- Bekci B, Cengiz C, Cengiz B 2012. Evaluating urban biodiversity in terms of user preferences: Urban residential landscape in Bartýn (Turkey). *Fresenius Environmental Bulletin*, 21: 1626-1635.
- Bekci B, Taskan G, Bogenc C 2013. A habitant-focused approach to the concept of eco-city at industrialized cities with a rural city character (Bartýn-Karabük) Turkey sample. *Journal of Food, Agriculture and Environment*, 11: 867-872.
- Belojevic G, Evans GE, Paunovic K, Jakovljevic B 2012. Traffic noise and executive functioning in urban primary school children: The moderating role of gender. *Journal of Environmental Psychology*, 32: 337-341.
- Bernatzky A 1978. *Tree Ecology and Preservation*. New York: Elsevier Scientific Publishing Company.
- Calixto A, Diniz FB, Zannin PHT 2003. The statistical modeling of road traffic noise in an urban setting. *Cities*, 20: 23-29.
- Cengiz C, Cengiz B, Bekci B 2012. Environmental quality analysis for sustainable urban public green space management in Bartın, Turkey. *Journal of Food, Agriculture and Environment*, 10: 938-946.
- Erdogan E, Yazgan ME 2009. Landscaping in reducing traffic noise problem in cities: Ankara case. *African Journal of Agricultural Research*, 4: 1015-1022.
- Fang CF, Ling DL 2003. Investigation of the noise reduction provided by tree belts. *Landscape and Urban Planning*, 63: 187-195.
- Fang CH, Ling DL 2005. Guidance for noise reduction provided by tree belts. *Landscape and Urban Planning*, 11: 29-34.
- Fritschi L, Brown L, Kim R, Schwela D, Kephelopoulou S 2011. *Burden of Disease From Environmental Noise – Quantification of Healthy Life Years Lost in Europe*. Bonn: WHO European Centre for Environment and Health.
- Haq I, Hussain T, Farooq H, Ahmad MR 2014. Evaluation of the traffic noise pollution at some busiest sites of Faisalabad City, Pakistan. *Academic Research International*, 5: 23-26.
- Huddart L 1990. *The Use of Vegetation for Traffic Noise Screening*. Berkshire: Transport Research Laboratory.
- Hunashal RB, Patil YB 2012. Assessment of noise pollution indices in the city of Kolhapur, India. *Procedia - Social and Behavioral Sciences*, 37: 448-457.
- Islam MN, Rahman KS, Bahar MM, Habib MA, Ando K et al. 2012. Pollution attenuation by roadside greenbelt in and around urban areas. *Urban Forestry and Urban Greening*, 11: 460-464.
- Kelkit A 2003. Environmental problems of Canakkale City and solutions. *International Journal of Environment and Pollution*, 19: 66-74.
- Kempen EE, Kamp I, Stellato RK, Lopez-Barrio I, Haines MM et al. 2009. Children's annoyance reactions to aircraft and road traffic noise. *Journal of Acoustical Society of America*, 125: 895-904.
- King RP, Davis JR 2003. Community noise: Health effects and management. *International Journal Hygiene Environmental Health*, 206: 123-131.
- Lam KC, Ng SL, Hui WC, Chan PK 2005. Environmental quality of urban parks and open spaces in Hong Kong. *Environmental Monitoring and Assessment*, 111: 55-73.

- Li B, Tao S, Dawson RW 2002. Evaluation and analysis of traffic noise from the main urban roads in Beijing. *Applied Acoustic*, 63: 1137-1142.
- Liu C, Fuertes E, Tiesler CMT, Birk M, Babisch W et al. 2014. The associations between traffic-related air pollution and noise with blood pressure in children: Results from the GINIplus and LISAPlus studies. *International Journal of Hygiene and Environmental Health*, 217: 499-505.
- Mansouri N, Pourmahabadian M, Ghasemkhani M 2006. Road traffic noise in down town area of Tahrán. *Iran Journal of Environmental Health Science and Engineering*, 3: 267-273.
- Mariscal-Ramires JA, Fernandes-Prieto JA, Canada-Bago J, Gadeo-Martos MA 2014. A new algorithm to monitor noise pollution adapted to resource-constrained devices. *Multimedia Tools and Applications*, 70: 1-15.
- Miedema H 2007. Annoyance caused by environmental noise: Elements for evidence-based noise policies. *Journal of Social Issues*, 63: 41-57.
- Mitra A 2008. Diabetes and stress: A Review. *Studies on Ethno-Medicine*, 2: 131-135.
- Morillas JM, Escobar VG, Sierra JAM, Gomez RV, Carmona JT 2002. An environmental study in the city of Cáceres (Spain). *Applied Acoustics*, 63: 1061-1070.
- Moudon AV 2009. Real noise from the urban environment how ambient community noise affects health and what can be done about it. *American Journal of Preventive Medicine*, 37: 167-171.
- Mukhola MS 2014. Street-food vending: Training directed at better food handling and associated environmental issues. *Anthropologist*, 17: 251-258.
- Noori K, Zand F 2013. An investigation of traffic noise pollution effects of citizens' general and mental health (Case study: Kermanshah City). *Journal of Novel Applied Science*, 2: 344-349.
- Onder S, Kocbeker Z 2012. Importance of the green belts to reduce noise pollution and determination of roadside noise reduction effectiveness of bushes in Konya, Turkey. *International Journal of Agricultural, Biosystems Science and Engineering*, 6: 26-29.
- Oyedepo OS, Abdullahi AS 2009. A comparative study of noise pollution levels in some selected areas in Ibadan metropolis, Nigeria. *Environmental Monitoring and Assessment*, 120: 499-525.
- Ozbilen A, Var M 1992. Gürültü kirliliğinin doğal elemanlarla çözümlenmesi için Doğu Karadeniz Bölgesinde gürültü kirliliğine karşı etkin olan doğal eleman türleriyle Trabzon'da bir örnek çözüm. *Ekoloji Çevre Dergisi*, 1: 17-22.
- Ozdemir B, Demirel Ö, Yasar Y, Sarýkoc E, Pirselimoglu Z 2008. An Approach of Solution to the Problem of Noise Pollution at the City Parks Located Among the Coastal Highway in Trabzon By Using Plant Material. *Paper Presented in International Conference on Ecological Protection of Planet Earth in Karadeniz Technical University, Trabzon*, June 12 to 15, 2008.
- Ozer S, Irmak MA, Yilmaz H 2008. Determination of roadside noise reduction effectiveness of *Pinus sylvestris* L. and *Populus nigra* L. in Erzurum, Turkey. *Environment Monitoring Assessment*, 144: 191-197.
- Panadya GH 2003. Assessment of traffic noise and its impact on the community. *International Journal of Environmental Studies*, 60: 595-602.
- Pathak V, Tripathi BD, Mishra VK 2008. Dynamics of traffic noise in a tropical city Varanasi and its abatement through vegetation. *Environmental Monitoring and Assessment*, 146: 67-75.
- Paunovic K, Belojevic G, Jakovljevic B, Stojanov V, Ilic ZJ 2009. The effects of traffic noise on blood pressure of children aged 7-11 in Belgrade. *Acustica*, 95: 47-48.
- Ralte L, Ralte L, Lalramnghinglova H 2013. Assessment on different levels of noise pollution in Aizawl City, Mizoram, India. *Science Vision*, 13: 157-161.
- Renterghem TV, Botteldooren D, Verheyen K 2012. Road traffic noise shielding by vegetation belts of limited depth. *Journal of Sound and Vibration*, 331: 2404-2425.
- Renterghem TV 2014. Guidelines for optimizing road traffic noise shielding by non-deep tree belts. *Ecological Engineering*, 69: 276-286.
- Renterghem TV, Attenborough K, Maennel M, De-france J, Horoshenkov K et al. 2014. Measured light vehicle noise reduction by hedges. *Applied Acoustics*, 78: 19-27.
- Samara Th, Tsitsoni Th 2007. Road Traffic Noise Reduction by Vegetation in the Ring Road of a Big City. *Paper Presented in the International Conference on Environmental Management, Engineering, Planning and Economics in Skiathos Palace Hotel, Skiathos*, June 24 to 28, 2007.
- Subramani T, Kavitha M, Sivaraj KP 2012. Modeling of traffic noise pollution. *International Journal of Engineering Research and Applications*, 2: 3175-3182.
- Sukru D, Celalettin O, Hakan K, Sahin K 2006. Noise pollution and map Konya city in Turkey. *Journal of International Environmental Application and Science*, 1: 63-72.
- Tsitsoni Th, Batala E, Zagas Th 2005. Management of Urban and Suggestions for its Upgrade in the Municipality of Thessaloniki. *Paper Presented in the 12th Panhellenic Conference of the Scientific Society on Forest Science in Aristotle University of Thessaloniki, Drama*, October 2 to 4, 2005.
- Tyagi V, Kumar K, Jain VK 2006. A Study of the spectral characteristics of traffic noise attenuation by vegetation belts in Delhi. *Applied Acoustics*, 67: 926-935.
- Vasilakopoulou K, Kolokotsa D, Santamouris M 2014. Urban heat island mitigation techniques for sustainable cities. In: Stamatina Th Rassia, Panos M Pardalos (Eds.): *Cities for Smart Environmental and Energy Futures*. Berlin: Springer, pp. 215-233.
- WHO 1993. The Environmental Health Criteria Document on Community Noise. *Report EUR/HFA Target 24*. Copenhagen: WHO.
- WHO 2000. *Guidelines for Community Noise*. Geneva: WHO.
- WHO 2011. *Burden of Disease from Environmental Noise – Quantification of Healthy Life Years Lost in Europe*. Copenhagen: WHO Regional Office for Europe.
- Yang F, Bao ZY, Zhu ZJ 2011. An assessment of psychological noise reduction by landscape plants. *International Journal of Environmental Research and Public Health*, 8: 1032-1048.
- Zannin PT, Szeremetta B 2003. Evaluation of noise pollution in the botanical garden in Curitiba, Parana, Brazil. *Public Health Reports*, 19: 683-686.
- Zannin PHT, Ferreira AMC, Szeremetta B 2006. Evaluation of noise pollution urban parks. *Environmental Monitoring and Assessment*, 118: 423-433.