

Effects of Warm-up Exercises on Spirometric Measurements in Athletes and Sedentary Individuals

Aynur Otag¹ and Ilhan Otag²

*¹Department of Physiotherapy and Rehabilitation, Health Sciences Faculty,
Cumhuriyet University, Sivas, Turkey*

²Vocational School of Health, Cumhuriyet University, Sivas, Turkey

KEYWORDS Athlete. Exercise. Sedentary. Stretching. Spirometry

ABSTRACT Regular physical activity is important for health. The present study aimed to examine the effects of warm-up exercises involving stretching on Spiro metric measurements in 30 athletes and 30 sedentary individuals. This study was designed as a randomized, crossover study. Initially, Spiro metric measurements were conducted before and after treadmill exercise in accordance with the Bruce protocol. After 15 days, Spirometric measurements were conducted before and after a warm-up program and the same exercise test. With warm-up exercises, the forced expiratory volume in 1 second increased in the athlete group. Additionally, the peak expiratory flow, forced inspiratory flow (25-75%), and $V_{max_{25}}$ increased with warm-up exercises. However, warm-up exercises did not influence Spiro metric measurements in the sedentary group. Among athletes, ventilation volume values increased generally with warm-up exercises. For this reason, we can say that warm-up exercises increase muscle strength, resulting in optimization of aerobic function.